

YAGAI BOUBA¹, ADAMOU SOULEYMANOU², AUDREY RAISSA DJOMO DZADDI¹, FATIMA MOULIOM NKAIN³, EBIAMA LIFANDA¹, EDWIGE OMONA², INGRID KENKO², ANTOINE SOCPA², MICHELINE MARIE JOSE ESSI², HADJA HAMSATOU CHERIF¹, SERGE CLOTAIRE BILLONG¹, ANNE CECILE ZOUNG-KANYI³

1 Central Technical Group, National AIDS Control Committee, Yaoundé, Cameroon; 2 Division of Health Operational Research, Ministry of Public Health, Yaoundé, Cameroon; 3 University of Yaoundé 1, Yaoundé, Cameroon

BACKGROUND

- In Cameroon, some key populations such as adolescents, young people, men and sex workers are less reached by the routine screening services.
- To reach the global and national testing goals, innovative strategies such as HIV self testing (HIVST) might be needed.
- Cameroon has recently adopted a guidelines for the implementation of HIVST.
- The UNITAID-funded HIV Self-Testing Africa (STAR) Initiative, which aim to provide a solid foundation for the introduction and rapid scale-up of HIVST based on evidence has been recently implemented in several countries, including Cameroon.

In Cameroon, evidence from the STAR Initiative showed that oral HIV self testing is an effective approach to detect new HIV infections among the populations which are hard to reach with the routine testing mechanisms; especially in the models considering the partners of people living with HIV/AIDS and HIV testing sites.

On the other hand, the community and workplace distribution models of oral HIV self testing need to be adjusted.




Figure 1. Regression model of factors associated with positivity among self testers

Variables	Univariate model		Multivariate Model	
	OR (95% CI)	P-value	aOR (95% CI)	P-value
Region				
Centre	1		1	
Littoral	0.303 (0.210-0.438)	<0.001	0.494 (0.267-0.914)	0.025
Sud	0.929 (0.750-1.149)	0.496	0.683 (0.543-0.860)	0.001
Distribution type				
Primary	1		1	
Secondary	10.041 (7.995-12.609)	<0.001	0.359 (0.232-0.556)	<0.001
Distribution models				
ANC/PNC/MCHC	1		1	
Partners of PLWHA	3.761 (3.080-4.593)	<0.001	4.007 (3.265-4.917)	<0.001
Workplace	0.272 (0.190-0.387)	<0.001	0.176 (0.089-0.349)	<0.001
Community	0.075 (0.049-0.116)	<0.001	0.033 (0.017-0.064)	<0.001
HTS	3.791 (1.953-7.357)	<0.001	2.906 (1.457-5.794)	0.002
Age categories				
<25	0.142 (0.107-0.187)	<0.001	0.683 (0.490-0.952)	0.024
25-39	0.935 (0.769-1.137)	0.501	0.903 (0.739-1.102)	0.315
≥40	1		1	
Sex				
Male	1		1	
Female	1.189 (0.989-1.429)	0.065		
HIV testing history				
At least one	1		1	
Never	0.814 (0.635-1.042)	0.102	1.702 (1.312-2.208)	<0.001
Unknown	2.213 (1.036-4.728)	0.040	2.604 (1.181-5.740)	0.018

ANC: antenatal clinic, CI: confidence interval, HTS: HIV testing site; MCHC: mother and child health clinic, OR: odds ratio, PLWHIV: people living with HIV, PNC: postnatal clinic. All factors with a p-value <0.005 was adjusted in the multivariable analysis. The model was adjusted for region, distribution type, distribution models, age categories, and HIV testing history.

CONCLUSIONS

HIVST using M1, M2 & M5 distribution models is an effective testing strategy in Cameroon.

Funding: UNITAID funded HIV Self-Testing Africa (STAR) Initiative

Correspondence: Yagai Bouba, National AIDS Control Committee (NACC), Yaoundé, Cameroon; romeobouba@yahoo.fr

RESULTS

Table 1. Outcome of HIV testing by self-tester according to sex, age, region, and distribution models.

Variable	Overall	Non reactive	Reactive	Invalid	P-value
Region, n (%)					
Centre	23008	22357 (97.2)	569 (2.5)	82 (0.4)	
Littoral	5920	5848 (98.8)	69 (1.2)	3 (0.1)	<0.001
South	7006	6790 (96.9)	187 (2.7)	29 (0.4)	
Distribution type, n (%)					
Primary	23854	23534 (98.7)	218 (0.9)	102 (0.4)	
Secondary	12080	11461 (94.9)	607 (5.0)	12 (0.1)	<0.001
Distribution models, n (%)					
ANC	7630	7376 (96.7)	246 (3.2)	8 (0.1)	
Partners of PVVH	4539	4157 (91.6)	378 (8.3)	4 (0.1)	
Workplace	7309	7218 (98.8)	85 (1.2)	6 (0.1)	<0.001
Community	16309	16111 (98.8)	102 (0.6)	96 (0.6)	
HTS	147	133 (90.5)	14 (9.5)	0 (0.0)	
Age categories, years, n (%)					
<25	18415	18142 (98.5)	175 (1.0)	98 (0.5)	
25-39	11689	11252 (96.3)	424 (3.6)	13 (0.1)	<0.001
≥40	5830	5601 (96.1)	226 (3.9)	3 (0.1)	
Sex, n (%)					
Male	25710	25064 (97.5)	570 (2.2)	76 (0.3)	0.144
Female	10224	9931 (97.1)	255 (2.5)	38 (0.4)	
HIV testing History, n (%)					
At least one	29703	28949 (97.5)	679 (2.3)	75 (0.3)	
Never	6019	5843 (97.1)	137 (2.3)	39 (0.6)	<0.001
Unknown	212	203 (95.8)	9 (4.2)	0 (0.0)	
Total	35934	34995 (97.4)	825 (2.3)	114 (0.3)	

ANC: antenatal clinic, CI: confidence interval, HTS: HIV testing site, MCHC: mother and child health clinic, OR: odds ratio, PLWHIV: people living with HIV, PNC: postnatal clinic. P-values were calculated using the Chi-square test. P-values underlined in bold are those which are significant, with statistically significant level fixed at $p < 0.05$.

OBJECTIVE

To evaluate the effectiveness of HIVST in Cameroon and identify predictors of HIV seropositivity among self-testers.

METHODS

- The study was conducted from 2021-2022 in the Centre, Littoral & Southern regions of Cameroon;
- Eligibility criteria for the HIVST distribution were: (i) ≥18 years old; (ii) Men at risk; (iii) partners of HIV+ people; and (iv) Youths 18 to 24 years old at risk;
- The HIVST kits were distributed either directly or indirectly according to 5 models: (i) Antenatal, postnatal, and maternal and child clinics (ANC/PNC/MCH) (M1); (ii) Partners of other HIV+ (M2); (iii) Workplace (M3); (iv) Community (M4); and (v) HIV testing services (HTS) (M5);
- HIVST was performed on oral fluid with the OraQuick® HIV-1/2 test;
- Reactive and invalid test results from the HIVST were confirmed following the national algorithm at a health facility;
- Regression model was performed to identify factors associated with positivity among HIV self testers.